arXiv:0907.1505v1 [math.AT] 9 Jul 2009

(NON-)KOSZULITY OF OPERADS FOR n-ARY ALGEBRAS, COHOMOLOGY AND DEFORMATIONS

MARTIN MARKL AND ELISABETH REMM

ABSTRACT. We investigate Koszulity of operads for various n-ary algebras. We then focus to algebras with one anti-associative operation. Since the corresponding operad is not Koszul, the deformation cohomology differs from the standard one. We describe the relevant part of the deformation cohomology for this type of algebras using the minimal model for the anti-associative operad. In the remaining sections we discuss free partially associative algebras and formulate open problems.

CONTENTS

1. Introduction 1
2. Duality for quadratic operads revisited 2
3. Four families of n-ary algebras 4
4. Koszulity - the case study 6
5. Cohomology of algebras over non-Koszul operads – an example 11
6. Free partially associative n-algebras 16
7. Open problems 19
References 20

1. INTRODUCTION

We study Koszulity of operads for various n-ary algebras, i.e. algebras with an n-multilinear operation satisfying a specific version of associativity. We will then focus to the particular case of algebras with one anti-associative operation, i.e. an operation $a, b \mapsto ab$ satisfying $a(bc) + a(bc) = 0$ for each a, b and c. The corresponding operad $\widetilde{\mathcal{A}ss}$ is not Koszul, so the deformation cohomology differs from the "standard" one. We describe the standard cohomology for this type of algebras and the relevant part of the deformation cohomology based on the minimal model of $\widetilde{\mathcal{A}ss}$. We also give a description of free partially associative algebras which, in the Koszul cases, coincides with the one given in [7].

The first author was supported by the grant GA ČR 201/08/0397 and by the Academy of Sciences of the Czech Republic, Institutional Research Plan No. AV0Z10190503.

We are aware that most of the results were known or anticipated by specialists. Let us mention at least the work of H. Ataguema and A. Makhlouf [1], V. Dotsenko and A. Khoroshkin [2], A.V. Gnedbaye [7], E. Hoffbeck [10] and the talk given by J.-L. Loday at the Winter School in Srní, in January 2008.

The aim of this paper is to present these scattered results in a systematic way and explain some common misconceptions related to Koszul duals of operads which are quadratic but not binary, see Remark 2. The novel part is a description of the deformation cohomology for anti-associative algebras. To our knowledge, this paper is the first place where some nontrivial information about the minimal model of a non-Koszul operad and the related cohomology appears.

Conventions. The basic reference for operads, quadratic duality an Koszulness is [6], our notation and terminology will also be based on [16] and [17]. We will work with operads in the category of chain complexes over a field $\mathbf{k}$ of characteristic zero though, in the light of [5], most if not all results remain valid over the ring of integers.

2. Duality for quadratic operads revisited

All ideas recalled in this section are already present in [6], but we want to emphasize some specific features of the non-binary case related to a degree shift in the Koszul dual which seem to be overlooked, see Remark 2 below.

Fix a natural $n \geq 2$ and assume $E = \{E(a)\}_{a\geq 2}$ is a Σ-module such that $E(a) = 0$ if $a \neq n$. We will study operads $\mathcal{P}$ of the form $\mathcal{P} = \Gamma(E)/(R)$, where $\Gamma(E)$ is the free operad generated by E and (R) the operadic ideal generated by a subspace $R \subset \Gamma(E)(2n-1)$. Operads of this type are called *quadratic*, or *binary* quadratic if $n = 2$.

Let $E^\vee = \{E^\vee(a)\}_{a\geq 2}$ be a Σ-module with

$$E^\vee(a) := \begin{cases} \mathrm{sgn}_a \otimes \uparrow^{a-2} E(a)^\#, & \text{if } a = n \text{ and} \\ 0, & \text{otherwise} \end{cases}$$

where $\uparrow^{a-2}$ denotes the suspension iterated $a-2$ times, sgn_a the signum representation of the symmetric group Σ_a, and $\#$ the linear dual of a graded vector space with the induced representation. Recall that $V^\# := \mathit{Hom}(V, \mathbf{k})$, so $(V^\#)_d = (V_{-d})^\#$. There is a non-degenerate, Σ_{2n-1}-equivariant pairing

$$\langle - | - \rangle : \Gamma(E^\vee)(2n-1) \otimes \Gamma(E)(2n-1) \to \mathbf{k} \tag{1}$$

determined by requiring that

$$\langle \uparrow^{n-2} e' \circ_i \uparrow^{n-2} f' \mid e'' \circ_j f'' \rangle := \delta_{ij}(-1)^{(i+1)(n+1)} e'(e'') f'(f'') \in \mathbf{k},$$

for arbitrary $e', f' \in E(n)^\#$, $e'', f'' \in E(n)$.

Definition 1. *The* Koszul *or* quadratic *dual of the quadratic operad* $\mathcal{P} = \Gamma(E)/(R)$ *as above is the quotient*

$$\mathcal{P}^! := \Gamma(E^\vee)/(R^\perp),$$

where $R^\perp \subset \Gamma(E^\vee)(2n-1)$ *is the annihilator of* $R \subset \Gamma(E)(2n-1)$ *in the pairing (1), and* $(R^\perp)$ *the operadic ideal generated by* $R^\perp$.

Remark 2. If $\mathcal{P}$ is a quadratic operad generated by an operation of arity n and degree d, then the generating operation of $\mathcal{P}^!$ has the same arity but degree $-d+n-2$, i.e. for $n \neq 2$ (the non-binary case) the Koszul duality *may not* preserve the degree of the generating operation! There seemed to be a common misunderstanding that the Koszul duality preserves the degree of generating operations even in the non-binary case. This resulted in several wrong statements about the Koszulity of n-ary algebras.

Recall [17, Definition II.3.15] that the *operadic suspension* $\mathbf{s}\mathcal{P}$ of an operad $\mathcal{P} = \{\mathcal{P}(a)\}_{a\geq 1}$ is the operad $\mathbf{s}\mathcal{P} = \{\mathbf{s}\mathcal{P}(a)\}_{a\geq 1}$, where $\mathbf{s}\mathcal{P}(a) := \mathrm{sgn}_a \otimes \uparrow^{a-1} \mathcal{P}(a)$, with the structure operations induced by those of $\mathcal{P}$ in the obvious way taking into account, as always, the Koszul sign rule requiring that whenever we interchange two "things" of odd degrees, we multiply the sign by -1. The *cooperadic suspension* $\mathbf{s}\mathcal{C}$ of a cooperad $\mathcal{C}$ is defined analogously.

The *cobar construction* [17, Definition II.3.9] of a coaugmented cooperad $\mathcal{C}$ is a dg-operad $\Omega(\mathcal{C})$ of the form $\Omega(\mathcal{C}) = (\Gamma(\downarrow \overline{\mathbf{s}\mathcal{C}}), \partial_\Omega)$. Here $\mathbf{s}$ denotes the cooperadic suspension recalled above, $\overline{\mathbf{s}\mathcal{C}}$ the coaugmentation coideal of the coaugmented cooperad $\mathbf{s}\mathcal{C}$, and $\downarrow$ the component-wise desuspension. The differential ∂_Ω is induced by the structure operations of the cooperad $\mathcal{C}$. If $\mathcal{P} = \{\mathcal{P}(a)\}_{a\geq 1}$ is an augmented operad with finite-dimensional components, the component-wise linear dual $\mathcal{P}^\# = \{\mathcal{P}(a)^\#\}_{a\geq 1}$ is a coaugmented cooperad. The composition $\mathsf{D}(\mathcal{P}) := \Omega(\mathcal{P}^\#)$ of the linear dual with the cobar construction is the *dual bar construction* of $\mathcal{P}$. For $\mathcal{P}$ quadratic, there exist a natural map $\mathsf{D}(\mathcal{P}^!) \to \mathcal{P}$ of dg-operads. Let us recall the central notion of [6].

Definition 3. *A quadratic operad* $\mathcal{P}$ *is* Koszul *if the natural map* $\mathsf{D}(\mathcal{P}^!) \to \mathcal{P}$ *is a homology equivalence.*

The notation and terminology concerning the (co)bar construction has not been standardized. For instance, the use of the (co)operadic suspension $\mathbf{s}$ in our definition of $\Omega(\mathcal{C})$ is purely conventional and guarantees that $\Omega(-)$ does not change the degrees of elements of arity 2. In recent literature, the Koszul dual is sometimes defined as a functor $\mathcal{P} \mapsto \mathcal{P}^{¡}$ from the category of quadratic operads to the category of quadratic *co*operads. Our $\mathcal{P}^!$ recalled above is then the composition of this co-operadic Koszul dual with the component-wise linear dual, $\mathcal{P}^! = (\mathcal{P}^{¡})^\#$. In our choice of conventions we, however, tried to stay as close to [6] as possible.

3. Four families of n-ary algebras

We introduce four families of quadratic operads and describe their Koszul duals. These families cover all examples of 'n-ary algebras' with one operation without symmetry which we were able to find in the literature.

Let V be a graded vector space, $n \geq 2$, and $\mu : V^{\otimes n} \to V$ a degree d multilinear operation symbolized by



.

We say that $A = (V, \mu)$ is a *degree d totally associative n-ary algebra* if, for each $1 \leq i, j \leq n$,

$$\mu\left(1\!\!1^{\otimes i-1} \otimes \mu \otimes 1\!\!1^{\otimes n-i}\right) = \mu\left(1\!\!1^{\otimes j-1} \otimes \mu \otimes 1\!\!1^{\otimes n-j}\right),$$

where $1\!\!1 : V \to V$ is the identity map. Graphically, we demand that



for each i, j for which the above compositions make sense. Observe that degree 0 totally associative 2-algebras are ordinary associative algebras.

In the following definitions, $\Gamma(\mu)$ will denote the free operad on the Σ-module E_μ with

$$E_\mu(a) = \begin{cases} \text{the regular representation } \mathbf{k}[\Sigma_n] \text{ generated by } \mu, & \text{if } a = n \text{ and} \\ 0, & \text{otherwise.} \end{cases}$$

Definition 4. *We denote $t\mathcal{A}ss^n_d$ the operad for totally associative n-ary algebras with operation in degree d, that is,*

$$t\mathcal{A}ss^n_d := \Gamma(\mu)/(R_{t\mathcal{A}ss^n_d})$$

with μ an arity n generator of degree d and

$$R_{t\mathcal{A}ss^n_d} := Span\left\{\mu \circ_i \mu - \mu \circ_j \mu, \text{ for } i, j = 1, \ldots, n\right\}.$$

We call $A = (V, \mu)$ a *degree d partially associative n-ary algebra* if the following single axiom is satisfied:

$$\sum_{i=1}^{n} (-1)^{(i+1)(n-1)} \mu\left(1\!\!1^{\otimes i-1} \otimes \mu \otimes 1\!\!1^{\otimes n-i}\right) = 0. \tag{2}$$

Degree 0 partially associative 2-ary algebras are classical associative algebras. More interesting observation is that degree $(n-2)$ partially associative n-ary algebras are the same as A_∞-algebras $A = (V, \mu_1, \mu_2, \ldots)$ [11, §1.4] which are "meager" in that they satisfy $\mu_k = 0$ for $k \neq n$. Symmetrizations of these *meager A_∞-algebras* are *Lie n-algebras* in the sense of [9].

Definition 5. *We denote $p\mathcal{A}ss^n_d$ the operad for partially associative n-ary algebras with operation in degree d. Explicitly,*

$$p\mathcal{A}ss^n_d := \Gamma(\mu)/\Big(\sum_{i=1}^{n}(-1)^{(i+1)(n-1)}\mu\circ_i\mu\Big)$$

with μ a generator of degree d and arity n.

It follows from the above remarks that $t\mathcal{A}ss^2_0 = p\mathcal{A}ss^2_0 = \mathcal{A}ss$, where $\mathcal{A}ss$ is the operad for associative algebras. We are going to introduce the remaining two families of operads. Recall that $\mathbf{s}$ denotes the operadic suspension and $\mathbf{s}^{-1}$ the obvious inverse operation.

Definition 6. *We define $\widetilde{t\mathcal{A}ss}^n_d := \mathbf{s}\, t\mathcal{A}ss^n_{d-n+1}$ and $\widetilde{p\mathcal{A}ss}^n_d := \mathbf{s}^{-1}p\mathcal{A}ss^n_{d+n-1}$.*

We leave as an exercise to verify that $\widetilde{t\mathcal{A}ss}^n_d$-algebras are structures $A = (V,\mu)$, where $\mu : V^{\otimes n}\to V$ is a degree d linear map satisfying, for each $1\leq i,j\leq n$,

$$(-1)^{i(n+1)}\mu\left(1\!\!1^{\otimes i-1}\otimes\mu\otimes 1\!\!1^{\otimes n-i}\right) = (-1)^{j(n+1)}\mu\left(1\!\!1^{\otimes j-1}\otimes\mu\otimes 1\!\!1^{\otimes n-j}\right).$$

Likewise, $\widetilde{p\mathcal{A}ss}^n_d$-algebras are similar structures, but this time satisfying

$$\sum_{i=1}^{n}\mu\left(1\!\!1^{\otimes i-1}\otimes\mu\otimes 1\!\!1^{\otimes n-i}\right) = 0.$$

Definition 7. *Let $\widetilde{\mathcal{A}ss} := \widetilde{t\mathcal{A}ss}^2_0 = \widetilde{p\mathcal{A}ss}^2_0$. Explicitly, $\widetilde{\mathcal{A}ss}$-algebras are structures $A = (V,\mu)$ with a degree 0 bilinear operation $\mu : V\otimes V\to V$ satisfying*

$$\mu(\mu\otimes 1\!\!1) + \mu(1\!\!1\otimes\mu) = 0$$

or, in elements

$$a(bc) + (ab)c = 0, \tag{3}$$

for $a,b,c\in V$. We call these objects anti-associative *algebras.*

Anti-associative algebras can be viewed as associative algebras with the associativity taken with the opposite sign which explains their name. Similarly, $t\mathcal{A}ss^2_1 = p\mathcal{A}ss^2_1$-algebras are associative algebras with operation of degree 1. The corresponding, essentially equivalent, operads are the simplest examples of non-Koszul operads, as we will see in Section 4. The proof of the following proposition is an exercise.

Proposition 8. *For each $n\geq 2$ and d, $(t\mathcal{A}ss^n_d)^! = p\mathcal{A}ss^n_{-d+n-2}$, $(p\mathcal{A}ss^n_d)^! = t\mathcal{A}ss^n_{-d+n-2}$, $(\widetilde{t\mathcal{A}ss}^n_d)^! = \widetilde{p\mathcal{A}ss}^n_{-d+n-2}$ and $(\widetilde{p\mathcal{A}ss}^n_d)^! = \widetilde{t\mathcal{A}ss}^n_{-d+n-2}$,*

$t\mathcal{A}ss^n_d$	n even	n odd
d even	yes	yes
d odd	no	no

$p\mathcal{A}ss^n_d$	n even	n odd
d even	yes	no
d odd	no	yes

$t\widetilde{\mathcal{A}ss}^n_d$	n even	n odd
d even	no	yes
d odd	yes	no

$p\widetilde{\mathcal{A}ss}^n_d$	n even	n odd
d even	no	no
d odd	yes	yes

FIGURE 1. Koszulness of the operads $t\mathcal{A}ss^n_d$, $p\mathcal{A}ss^n_d$, $t\widetilde{\mathcal{A}ss}^n_d$ and $p\widetilde{\mathcal{A}ss}^n_d$. "Yes" means that the corresponding operad is Koszul, "no" that it is not Koszul.

4. KOSZULITY - THE CASE STUDY

This section is devoted to the following statement organized in the tables of Figure 1. We, however, know to prove the non-Koszulity direction ("no" in Figure 1) only for $n \leq 7$ although we expect it to be true for arbitrary n, see also the notes following Theorem 9, Remark 18 and the first problem of Section 7.

Theorem 9. *The operad $t\mathcal{A}ss^n_d$ is Koszul if and only if d is even. The operad $p\mathcal{A}ss^n_d$ is Koszul if and only if n and d have the same parity. The operad $t\widetilde{\mathcal{A}ss}^n_d$ is Koszul if and only if n and d have different parities. The operad $p\widetilde{\mathcal{A}ss}^n_d$ is Koszul if and only if d is odd.*

The Koszulity part ("yes" in the tables of Figure 1) will follow from [10] and relations between the operads $t\mathcal{A}ss^n_d$, $p\mathcal{A}ss^n_d$, $t\widetilde{\mathcal{A}ss}^n_d$ and $p\widetilde{\mathcal{A}ss}^n_d$, see Proposition 11. The non-Koszulness part ("no" in Figure 1) will, for $n \leq 7$, follow in a similar fashion from Proposition 17. Since we do not know how to extend the proof of Proposition 17 for arbitrary n, the non-Koszulity part of Theorem 9 is, for $n \geq 8$, only conjectural.

In particular, the operads $\widetilde{\mathcal{A}ss}$ and $t\mathcal{A}ss^2_1 = p\mathcal{A}ss^2_1$ are not Koszul. Let us formulate useful

Lemma 10. *Let $\mathcal{P}^n_d$ be one of the operads above. Then $\mathcal{P}^n_d$ is Koszul if and only if $\mathcal{P}^n_{d+2}$ is Koszul, that is, only the parity of d matters.*

Proof. There is a 'twisted' isomorphism

$$\varphi : \mathcal{P}^n_d \xrightarrow{\cong} \mathcal{P}^n_{d+2}, \tag{4}$$

i.e. a sequence of equivariant isomorphisms $\varphi(a) : \mathcal{P}^n_d(a) \to \mathcal{P}^n_{d+2}(a)$, $a \geq 1$, that commute with the $\circ_i$-operations such that the component $\varphi(k(n-1)+1)$ is of degree $2k$, for $k \geq 0$.

To construct such an isomorphism, consider an operation μ' of arity n and degree d, and another operation μ'' of the same arity but of degree $d+i$. We leave as an exercise to verify that the assignment $\mu' \mapsto \mu''$ extends to a twisted isomorphism $\omega : \Gamma(\mu') \to \Gamma(\mu'')$ if and only if i is even.

Let $\mathcal{P}^n_d = \Gamma(\mu')/(R')$ and $\mathcal{P}^n_{d+2} = \Gamma(\mu'')/(R'')$. It is clear that the twisted isomorphism $\omega : \Gamma(\mu') \to \Gamma(\mu'')$ preserves the ideals of relations, so it induces a twisted isomorphism (4). A moment's reflection convinces one that φ induces similar twisted isomorphisms of the Koszul duals and the bar constructions. This, by Definition 3, gives the lemma. □

Proposition 11. *The operads marked "yes" in the tables of Figure 1 are Koszul.*

Proof. The operads $t\mathcal{A}ss^n_0$ are Koszul for all $n \geq 2$ by [10, § 7.2] (see also [7] for the case n even and $d = 0$). So, by Lemma 10, the operads $t\mathcal{A}ss^n_d$ are Koszul for all even d and $n \geq 2$, which gives the two "yes" in the upper left table of Figure 1.

The "yes" in the upper right table follow from the "yes" in the upper left table, the fact that an operad is Koszul if and only if its dual operad is Koszul, and the isomorphism $(p\mathcal{A}ss^n_d)^! = t\mathcal{A}ss^n_{-d+n-2}$ established in Proposition 8. The "yes" in the lower two tables in Figure 1 follow from the upper two tables and the fact that the suspension preserves Koszulity. □

Before we attend to the "no" part of Theorem 9, we recall an important criterion for Koszulity due to [6] which easily generalizes to the non-binary case. The *Poincaré* or *generating series* of a graded operad $\mathcal{P}_* = \{\mathcal{P}_*(a)\}_{a\geq 1}$ is defined by

$$g_{\mathcal{P}}(t) := \sum_{a\geq 1} \frac{1}{a!}\chi(\mathcal{P}(a))t^a,$$

where $\chi(\mathcal{P}(a))$ denotes the Euler characteristic of the graded vector space $\mathcal{P}_*(n)$,

$$\chi(\mathcal{P}(a)) := \sum_i (-1)^i \dim(\mathcal{P}_i(a)).$$

Observe that if $\mathcal{P}$ is not concentrated in degree zero, the Euler characteristic and thus also the Poincaré series may involve negative signs.

Theorem 12 ([6])**.** *If a quadratic operad $\mathcal{P}$ is Koszul, then its Poincaré series and the Poincaré series of its dual $\mathcal{P}^!$ are tied by the functional equation*

$$g_{\mathcal{P}}(-g_{\mathcal{P}^!}(-t)) = t. \tag{5}$$

Our first task will therefore be to describe the Poincaré series of the family $t\mathcal{A}ss^n_d$ which generates, via the duality and suspension, all the remaining operads.

Lemma 13. *The generating function for the operad $t\mathcal{A}ss^n_d$ is*

$$g_{t\mathcal{A}ss^n_d}(t) := \begin{cases} \dfrac{t}{1-t^{n-1}}, & \text{if } d \text{ is even, and} \\ t - t^n + t^{2n-1}, & \text{if } d \text{ is odd.} \end{cases} \tag{6}$$

Proof. The components of the operad $t\mathcal{A}ss^n_d$ are trivial in arities different from $k(n-1)+1$, $k \geq 0$. The piece $t\mathcal{A}ss^n_d(k(n-1)+1)$ is generated by all possible $\circ_i$-compositions involving k instances of the generating operation μ, modulo the relations

$$\mu \circ_i \mu - \mu \circ_j \mu, \text{ for } i, j = 1, \ldots, n \tag{7}$$

which enable one to replace each $\mu \circ_i \mu$, $2 \leq i \leq n$, by $\mu \circ_1 \circ\mu$.

If the degree d is *even*, the operad $t\mathcal{A}ss^n_d$ is evenly graded, so the associativity [13, p. 1473, Eqn. (1)] of the $\circ_i$-operations does not involve signs. Therefore an arbitrary $\circ_i$-composition of k instances of μ can be brought to the form

$$\eta_k := (\cdots((\mu \circ_1 \mu) \circ_1 \mu) \circ_1 \cdots) \circ_1 \mu.$$

We see that $t\mathcal{A}ss^n_d(k(n-1)+1)$ is spanned by the set $\{\eta_k \circ \sigma;\ \sigma \in \Sigma_{k(n-1)+1}\}$, so

$$\dim(t\mathcal{A}ss^n_d(k(n-1)+1)) = (k(n-1)+1)!$$

and, by definition,

$$g_{t\mathcal{A}ss^n_d}(t) = \sum_{k \geq 0} t^{k(n-1)+1} = \frac{t}{1-t^{n-1}},$$

which verifies the even case of (6).

The *odd* case is subtler since the associativity [13, p. 1473, Eqn. (1)] may involve nontrivial signs. As in the even case we calculate that

$$\dim(t\mathcal{A}ss^n_d(k(n-1)+1)) = (k(n-1)+1)! \ \text{ for } k = 0, 1, 2, \tag{8}$$

because these small arities do not require the associativity.

If $k \geq 3$, we can still to bring each $\circ_i$-composition of k instances of μ to the form of the 'canonical' generator η_k, but we may get a nontrivial sign which may moreover depend on the way we applied the associativity. Relation (7) implies that

$$(\mu \circ_1 \mu) \circ_1 \mu = (\mu \circ_n \mu) \circ_1 \mu \tag{9}$$

in $t\mathcal{A}ss^n_d(3n-2)$. Applying (7) and the associativity [13, p. 1473, Eqn. (1)] several times, we get that

$$\begin{aligned}(\mu \circ_1 \mu) \circ_1 \mu &= \mu \circ_1 (\mu \circ_1 \mu) = \mu \circ_1 (\mu \circ_n \mu) = (\mu \circ_1 \mu) \circ_n \mu = (\mu \circ_n \mu) \circ_n \mu \\ &= \mu \circ_n (\mu \circ_1 \mu) = \mu \circ_n (\mu \circ_n \mu) = (\mu \circ_n \mu) \circ_{2n-1} \mu \\ &= (\mu \circ_1 \mu) \circ_{2n-1} \mu.\end{aligned} \tag{10}$$

Since the degree of μ is odd, the first line of the associativity [13, p. 1473, Eqn. (1)] implies

$$(\mu \circ_1 \mu) \circ_{2n-1} \mu = -(\mu \circ_n \mu) \circ_1 \mu$$

therefore (9) and (10) combine into

$$(\mu \circ_1 \mu) \circ_1 \mu = -(\mu \circ_1 \mu) \circ_1 \mu.$$

This means that $(\mu \circ_1 \mu) \circ_1 \mu = 0$ so $t\mathcal{A}ss^n_d(3n-2) = 0$. Since $t\mathcal{A}ss^n_d(k(n-1)+1)$ is, for $k \geq 3$, generated by $t\mathcal{A}ss^n_d(3n-2)$, we conclude that $t\mathcal{A}ss^n_d(k(n-1)+1) = 0$ for $k \geq 3$ which, along with (8), verifies the odd case of (7). □

Remark 14. The Poincaré series of an operad $\mathcal{P}$ and its suspension $\mathbf{s}\mathcal{P}$ are related by $g_{\mathbf{s}\mathcal{P}}(t) = -g_{\mathcal{P}}(-t)$. Lemma 13 thus implies that the generating series of the operad $t\widetilde{\mathcal{A}ss}^n_d = \mathbf{s}t\mathcal{A}ss^n_{d-n+1}$ equals

$$g_{t\widetilde{\mathcal{A}ss}^n_d}(t) := \begin{cases} t + (-1)^d t^n + t^{2n-1}, & \text{if } n \text{ and } d \text{ have the same parity, and} \\ \dfrac{t}{1-(-1)^d t^{n-1}}, & \text{if } n \text{ and } d \text{ have different parities.} \end{cases}$$

We do not know explicit formulas for the Poincaré series of $p\mathcal{A}ss^n_d$ and $p\widetilde{\mathcal{A}ss}^n_d$ except in the case $n = 2$ when these operads coincide with the corresponding (anti)-associative operads.

Example 15. It easily follows from the above calculations that, for the anti-associative operad $\widetilde{\mathcal{A}ss}$, one has

$$\widetilde{\mathcal{A}ss}(1) \cong \mathbf{k},\ \widetilde{\mathcal{A}ss}(2) \cong \mathbf{k}[\Sigma_2] \text{ and } \widetilde{\mathcal{A}ss}(3) \cong \mathbf{k}[\Sigma_3],$$

while $\widetilde{\mathcal{A}ss}(a) = 0$ for $a \geq 4$.

Let us return to our task of proving the non-Koszulity of the "no" cases in the tables of Figure 1. Our strategy will be to interpret (5) as saying that $-g_{\mathcal{P}^!}(-t)$ is a formal inverse of $g_{\mathcal{P}}(t)$ at 0. Since $g'_{\mathcal{P}}(0) = 1$, this unique formal inverse exists. In the particular case of $\mathcal{P} = t\mathcal{A}ss^n_d$, with d odd, this means that $-g_{p\mathcal{A}ss^n_{-d+n-2}}(-t)$ should be compared to a formal inverse of $g_{t\mathcal{A}ss^n_d}(t) = t - t^n + t^{2n-1}$. A simple degree count shows that $g_{p\mathcal{A}ss^n_{-d+n-2}}(t)$ is of the form

$$\begin{cases} t - A_1 t^n + A_2 t^{2n-1} - A_3 t^{3n-2} + \cdots, & \text{for } n \text{ even and} \\ t + A_1 t^n + A_2 t^{2n-1} + A_3 t^{3n-2} + \cdots, & \text{for } n \text{ odd,} \end{cases}$$

for some *non-negative* integers $A_1, A_2, A_3, \ldots$, therefore $-g_{p\mathcal{A}ss^n_{-d+n-2}}(-t)$ is in both cases the formal power series

$$t + A_1 t^n + A_2 t^{2n-1} + A_3 t^{3n-2} + \cdots \tag{11}$$

with non-negative coefficients. If we show that the formal inverse of $t - t^n + t^{2n-1}$ is not of this form, by Theorem 12 the corresponding operad $t\mathcal{A}ss^n_d$ is not Koszul.

Example 16. The Poincaré series of the operad $t\mathcal{A}ss^2_1$ is, by Lemma 13,

$$g_{t\mathcal{A}ss^2_1}(t) = t - t^2 + t^3.$$

One can compute the formal inverse of this function as

$$t + t^2 + t^3 - 4t^5 - 14t^6 - 30t^7 - 33t^8 + 55t^9 + \cdots.$$

The presence of negative coefficients implies that the operad $t\mathcal{A}ss^2_1$ is not Koszul, neither is the anti-associative operad $\widetilde{\mathcal{A}ss} = t\widetilde{\mathcal{A}ss}^2_0 = \mathbf{s}^{-1}t\mathcal{A}ss^2_1$.

Likewise, the Poincaré series of the operad $t\mathcal{A}ss_1^3$ equals

$$g_{t\mathcal{A}ss_1^3}(t) = t - t^3 + t^5$$

and we computed, using `Matematica`, the initial part of the formal inverse as

$$t + t^3 + 2t^5 + 4t^7 + 5t^9 - 13t^{11} - 147t^{13} + \cdots$$

The existence of negative coefficients again implies that the operad $t\mathcal{A}ss_1^3$ is not Koszul. The formal inverse of

$$g_{t\mathcal{A}ss_1^4}(t) = t - t^4 + t^7$$

up to the first negative term is

$$t + t^4 + 3t^7 + 11t^{10} + 42t^{13} + 153t^{16} + 469t^{19} + 690t^{22} - 5967t^{25} + \cdots$$

so $t\mathcal{A}ss_1^4$ is not Koszul.

The complexity of the calculation of the relevant initial part of the inverse of $g_{t\mathcal{A}ss_1^n}(t) = t - t^n + t^{2n-1}$ grows rapidly with n. We have, however, the following:

Proposition 17. *For $n \leq 7$, the formal inverse of $t - t^n + t^{2n-1}$ has at least one negative coefficient. Therefore the operads $t\mathcal{A}ss_d^n$ for d odd and $n \leq 7$ are not Koszul.*

Proof. The function $g(z) := z - z^n + z^{2n-1}$ is analytic in the complex plane $\mathbb{C}$. Its analytic inverse $g^{-1}(z)$ is a not-necessarily single-valued analytic function defined outside the points in which the derivative $g'(z)$ vanishes. Let us denote by $\mathfrak{Z}$ the set of these points, i.e.

$$\mathfrak{Z} := \{z \in \mathbb{C};\ g'(z) = 0\}.$$

The key observation is that, for $n \leq 7$, the equation $g'(z) = 0$ has no real solutions, $\mathfrak{Z} \cap \mathbb{R} = \emptyset$. Indeed, one has to solve the equation

$$g'(z) = 1 - nz^{n-1} + (2n-1)z^{2n-2} = 0 \tag{12}$$

which, after the substitution $w := y^{n-1}$ leads to the quadratic equation

$$1 - nw + (2n-1)w^2 = 0$$

whose discriminant $n^2 - 8n + 4$ is, for $n \leq 7$, negative.

Let $f(z)$ be the power series representing the branch at 0 of $g^{-1}(z)$ such that $f(0) = 0$. It is clear that $f(t)$ is precisely the formal inverse of $g(t)$ at 0. Suppose that

$$f(z) = z + a_2z^2 + a_3z^3 + a_4z^4 + \cdots,$$

with all coefficients $a_2, a_3, a_4, \ldots$ non-negative real numbers. Since $\mathfrak{Z} \neq \emptyset$ and obviously $0 \notin \mathfrak{Z}$, the radius of convergence of $f(z)$ at 0, which equals the radius of the maximal circle centered at 0 whose interior does not contain points in $\mathfrak{Z}$, is some number r with $0 < r < \infty$. Let $\mathfrak{z} \in \mathfrak{Z}$ be such that $|\mathfrak{z}| = r$. Since all coefficients of the power series f are positive, we have

$$|f(\mathfrak{z})| \leq f(|\mathfrak{z}|) = f(r),$$

so the function $f(r)$ must have singularity at the *real* point $r \in \mathbb{R}$, i.e. $g'(z)$ must vanish at r. This contradicts the fact that $g'(z) = 0$ has no real solutions. $\square$

Remark 18. Equation (12) has, for $n = 8$, two real solutions, $\mathfrak{z}_1 = \sqrt[7]{1/3}$ and $\mathfrak{z}_2 = \sqrt[7]{1/5}$. This means that the inverse function of $z - z^n + z^{2n-1}$ has two positive real poles and the arguments used in our proof of Proposition 17 do not apply.

We verified Proposition 17 using `Matematica`. The first negative coefficient in the inverse of $t - t^n + t^{2n-1}$ was at the power t^{57} for $n = 5$, at t^{161} for $n = 6$, and at t^{1171} for $n = 7$. For $n = 8$ we did not find any negative term of degree less than 10 000. It is indeed possible that all coefficients of the inverse of $t - t^8 + t^{15}$ are positive.

Proposition 17 together with the fact that the suspension and the !-dual preserves Koszulity imply the "no" entries of the tables in Figure 1 for $n \leq 7$.

5. COHOMOLOGY OF ALGEBRAS OVER NON-KOSZUL OPERADS – AN EXAMPLE

In this section we study anti-associative algebras introduced in Definition 7, i.e. structures $A = (V, \mu)$ with a degree-0 bilinear anti-associative multiplication $\mu : V^{\otimes 2} \to V$. We describe the 'standard' cohomology $H^*_{\widetilde{\mathcal{A}ss}}(A; A)_{\rm st}$ of an anti-associative algebra A with coefficients in itself and compare it to the relevant part of the deformation cohomology $H^*_{\widetilde{\mathcal{A}ss}}(A; A)$ based on the minimal model of the anti-associative operad $\widetilde{\mathcal{A}ss}$. Since $\widetilde{\mathcal{A}ss}$ is, by Theorem 9, not Koszul, these two cohomologies differ. While the standard cohomology has no sensible meaning, the deformation cohomology coincides with the triple cohomology [3, 4] and governs deformations of anti-associative algebras.

It was explained at several places [12, 13, 14, 15] how a, not-necessarily acyclic, quasi-free resolution $(\mathcal{P}, \partial = 0) \xleftarrow{\rho} (\mathscr{R}, \partial)$ of an operad $\mathcal{P}$, which we assume for simplicity non-dg and concentrated in degree 0, determines a cohomology theory for $\mathcal{P}$-algebras with coefficients in itself. If $\mathcal{P}$ is quadratic and if we take as $(\mathscr{R}, \partial)$ the dual bar construction (recalled in Section 2) of the quadratic dual $\mathcal{P}^!$, we get the 'standard' cohomology $H^*_{\mathcal{P}}(A; A)_{\rm st}$ as the cohomology of the 'standard' cochain complex

$$C^1_{\mathcal{P}}(A; A)_{\rm st} \xrightarrow{\delta^1_{\rm st}} C^2_{\mathcal{P}}(A; A)_{\rm st} \xrightarrow{\delta^2_{\rm st}} C^3_{\mathcal{P}}(A; A)_{\rm st} \xrightarrow{\delta^3_{\rm st}} C^4_{\mathcal{P}}(A; A)_{\rm st} \xrightarrow{\delta^4_{\rm st}} \cdots$$

in which $C^p_{\mathcal{P}}(A; A)_{\rm st} := \mathit{Hom}(\mathcal{P}^!(p) \otimes_{\Sigma_p} V^{\otimes p}, V)$, $p \geq 1$, and the differential $\delta^*_{\rm st}$ is induced from the structure of $\mathcal{P}^!$ and A, see [4, Section 8] or [17, Definition II.3.99]. This type of (co)homology was considered in the seminal paper [6].

The deformation (also called, in [12], the *cotangent*) cohomology uses the minimal model of $\mathcal{P}$ in place of $(\mathscr{R}, \partial)$. Recall [13, p. 1479] that the *minimal model* of an operad $\mathcal{P}$ is a homology isomorphism

$$(\mathcal{P}, 0) \xleftarrow{\rho} (\Gamma(E), \partial)$$

of dg-operads such that the image of ∂ consists of decomposable elements of the free operad $\Gamma(E)$ (the minimality). It is known [17, Section II.3.10] that each operad with $\mathcal{P}(1) \cong \mathbf{k}$ admits a minimal model unique up to isomorphism. The *deformation cohomology* $H^*_{\mathcal{P}}(A;A)$ is the cohomology of the complex

$$C^1_{\mathcal{P}}(A;A) \xrightarrow{\delta^1} C^2_{\mathcal{P}}(A;A) \xrightarrow{\delta^2} C^3_{\mathcal{P}}(A;A) \xrightarrow{\delta^3} C^4_{\mathcal{P}}(A;A) \xrightarrow{\delta^4} \cdots$$

in which $C^1_{\mathcal{P}}(A;A) := \mathit{Hom}(V,V)$ and

$$C^p_{\mathcal{P}}(A;A) := \mathit{Hom}(\textstyle\bigoplus_{q\geq 2} E_{p-2}(q) \otimes_{\Sigma_q} V^{\otimes q}, V), \quad \text{for } p \geq 2.$$

The differential δ^* is defined by the formula which can be found in [14, Section 2] or in the introduction to [15]. If $\mathcal{P}$ is quadratic Koszul, the dual bar construction of $\mathcal{P}^!$ is, by [13, Proposition 2.6], isomorphic to the minimal model of $\mathcal{P}$, thus the standard and deformation cohomologies coincide, giving rise to the 'standard' constructions such as the Hochschild, Harrison or Chevalley-Eilenberg cohomology.

Neither $H^*_{\mathcal{P}}(A;A)_{\mathrm{st}}$ nor $H^*_{\mathcal{P}}(A;A)$ have the 0th term. A natural H^0 exists only for algebras for which the concept of unitality makes sense. This is not always the case. Assume, for example, that an anti-associative algebra $A = (V,\mu)$ has a unit, i.e. and element $1 \in V$ such that $1a = a1 = a$, for all $a \in V$. Then the anti-associativity (3) with $c = 1$ gives $ab + ab = 0$, so $ab = 0$ for each $a, b \in V$.

Let us describe the standard cohomology $H^*_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}}$ of an anti-associative algebra $A = (V,\mu)$. The operad $\widetilde{\mathcal{A}ss}$ is, by Proposition 8, self-dual and it follows from the description of $\widetilde{\mathcal{A}ss} = \widetilde{\mathcal{A}ss}^!$ given in Example 15 that $H^*_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}}$ is the cohomology of

$$C^1_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}} \xrightarrow{\delta^1_{\mathrm{st}}} C^2_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}} \xrightarrow{\delta^2_{\mathrm{st}}} C^3_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}} \xrightarrow{\delta^3_{\mathrm{st}}} 0 \xrightarrow{0} 0 \xrightarrow{0} \cdots$$

in which $C^p_{\widetilde{\mathcal{A}ss}}(A;A) := \mathit{Hom}(V^{\otimes p}, V)$ for $p = 1, 2, 3$, and all higher $C^p_{\widetilde{\mathcal{A}ss}}(A;A)$'s are trivial. The two nontrivial pieces of the differential are basically the Hochschild differentials with "wrong" signs of some terms:

$$\delta^1(\varphi)(a,b) := a\varphi(b) - \varphi(ab) + \varphi(a)b, \text{ and}$$
$$\delta^2(f)(a,b,c) := af(b,c) + f(ab,c) + f(a,bc) + f(a,b)c,$$

for $\varphi \in \mathit{Hom}(V,V)$, $f \in \mathit{Hom}(V^{\otimes 2}, V)$ and $a,b,c \in V$. We abbreviated $\mu(a,b) = ab$, $\mu(a,\varphi(b)) = a\varphi(b)$, &c. One sees, in particular, that $H^p_{\widetilde{\mathcal{A}ss}}(A;A)_{\mathrm{st}} = 0$ for $p \geq 4$.

Let us describe the relevant part of the deformation cohomology of A. It can be shown that $\widetilde{\mathcal{A}ss}$ has the minimal model

$$(\widetilde{\mathcal{A}ss}, 0) \xleftarrow{\rho} (\Gamma(E), \partial)$$

with the generating Σ-module $E = \{E(a)\}_{a\geq 2}$ such that

- $E(2)$ is generated by a degree 0 bilinear operation $\mu_2 : V \otimes V \to V$,
- $E(3)$ is generated by a degree 1 trilinear operation $\mu_3 : V^{\otimes 3} \to V$,

- $E(4) = 0$, and
- $E(5)$ is generated by four 5-linear degree 2 operations $\mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4 : V^{\otimes 5} \to V$,

so the minimal model of $\widetilde{\mathcal{A}ss}$ is of the form

$$(\widetilde{\mathcal{A}ss}, 0) \xleftarrow{\alpha} (\Gamma(\mu_2, \mu_3, \mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4, \ldots), \partial).$$

Notice the gap in the arity 4 generators! We do not know the exact form of the pieces $E(a)$, $a \geq 6$, of the generating Σ-module E, but we know that they do not contain elements of degrees ≤ 2. Their Euler characteristics can be read off from the inverse of the Poincaré series $g_{\widetilde{\mathcal{A}ss}}(t) = t + t^2 + t^3$, and one gets

$$\begin{aligned}
&\chi(E(2)) = 1,\ \chi(E(3)) = -1,\ \chi(E(4)) = 0,\ \chi(E(5)) = 4,\\
&\chi(E(6)) = -14,\ \chi(E(7)) = 30,\ \chi(E(8)) = -33,\ \chi(E(9)) = -55, \ldots
\end{aligned}$$

The differential ∂ of the relevant generators is given by:

$$\begin{aligned}
\partial(\mu_2) &:= 0,\\
\partial(\mu_3) &:= \mu_2 \circ_1 \mu_2 + \mu_2 \circ_2 \mu_2,\\
\partial(\mu_5^1) &:= (\mu_2 \circ_2 \mu_3) \circ_4 \mu_2 - (\mu_3 \circ_3 \mu_2) \circ_4 \mu_2 + (\mu_2 \circ_1 \mu_2) \circ_3 \mu_3 - (\mu_3 \circ_1 \mu_2) \circ_3 \mu_2\\
&\quad + (\mu_2 \circ_1 \mu_3) \circ_1 \mu_2 - (\mu_3 \circ_1 \mu_2) \circ_1 \mu_2 + (\mu_2 \circ_1 \mu_3) \circ_4 \mu_2 - (\mu_3 \circ_2 \mu_2) \circ_4 \mu_2,\\
\partial(\mu_5^2) &:= (\mu_3 \circ_1 \mu_2) \circ_1 \mu_2 - (\mu_2 \circ_1 \mu_3) \circ_1 \mu_2 + (\mu_2 \circ_1 \mu_3) \circ_3 \mu_2 - (\mu_3 \circ_2 \mu_2) \circ_3 \mu_2\\
&\quad + (\mu_2 \circ_2 \mu_3) \circ_3 \mu_2 - (\mu_3 \circ_3 \mu_2) \circ_3 \mu_2 + (\mu_2 \circ_1 \mu_2) \circ_3 \mu_3 - (\mu_3 \circ_1 \mu_2) \circ_4 \mu_2,\\
\partial(\mu_5^3) &:= (\mu_3 \circ_2 \mu_2) \circ_4 \mu_2 - (\mu_2 \circ_2 \mu_3) \circ_2 \mu_2 + (\mu_3 \circ_2 \mu_2) \circ_2 \mu_2 - (\mu_2 \circ_1 \mu_2) \circ_2 \mu_3\\
&\quad + (\mu_2 \circ_1 \mu_3) \circ_3 \mu_2 - (\mu_2 \circ_1 \mu_3) \circ_1 \mu_2 + (\mu_2 \circ_1 \mu_2) \circ_1 \mu_3 - (\mu_3 \circ_1 \mu_2) \circ_2 \mu_2, \text{ and}\\
\partial(\mu_5^4) &:= (\mu_3 \circ_1 \mu_2) \circ_3 \mu_2 - (\mu_3 \circ_3 \mu_2) \circ_3 \mu_2 + (\mu_2 \circ_2 \mu_3) \circ_3 \mu_2 - (\mu_3 \circ_2 \mu_2) \circ_3 \mu_2\\
&\quad + (\mu_2 \circ_1 \mu_2) \circ_2 \mu_3 - (\mu_2 \circ_1 \mu_3) \circ_2 \mu_2 + (\mu_2 \circ_1 \mu_2) \circ_1 \mu_3 - (\mu_2 \circ_1 \mu_3) \circ_1 \mu_2.
\end{aligned}$$

One can make the formulas clearer by using the nested bracket notation. For instance, μ_2 will be represented by $(\bullet\bullet)$, μ_3 by $(\bullet\bullet\bullet)$, μ_5^2 by $(\bullet\bullet\bullet\bullet\bullet)^2$, $\mu_3 \circ_2 \mu_2$ by $(\bullet(\bullet\bullet)\bullet)$, &c. With this shorthand, the formulas for the differential read

$$\begin{aligned}
\partial(\bullet\bullet) &:= 0,\\
\partial(\bullet\bullet\bullet) &:= ((\bullet\bullet)\bullet) + (\bullet(\bullet\bullet)),\\
\partial(\bullet\bullet\bullet\bullet)^1 &:= (\bullet(\bullet\bullet(\bullet\bullet))) - (\bullet\bullet(\bullet(\bullet\bullet))) + ((\bullet\bullet)(\bullet\bullet\bullet)) - ((\bullet\bullet)(\bullet\bullet)\bullet)\\
&\quad + (((\bullet\bullet)\bullet\bullet)\bullet) - (((\bullet\bullet)\bullet)\bullet\bullet) + ((\bullet\bullet\bullet)(\bullet\bullet)) - (\bullet(\bullet\bullet)(\bullet\bullet)),\\
\partial(\bullet\bullet\bullet\bullet)^2 &:= (((\bullet\bullet)\bullet)\bullet\bullet) - (((\bullet\bullet)\bullet\bullet)\bullet) + ((\bullet\bullet(\bullet\bullet))\bullet) - (\bullet(\bullet(\bullet\bullet))\bullet)\\
&\quad + (\bullet(\bullet(\bullet\bullet)\bullet)) - (\bullet\bullet((\bullet\bullet)\bullet)) + ((\bullet\bullet)(\bullet\bullet\bullet)) - ((\bullet\bullet)\bullet(\bullet\bullet)),\\
\partial(\bullet\bullet\bullet\bullet)^3 &:= (\bullet(\bullet\bullet)(\bullet\bullet)) - (\bullet((\bullet\bullet)\bullet\bullet)) + (\bullet((\bullet\bullet)\bullet)\bullet) - ((\bullet(\bullet\bullet\bullet))\bullet)\\
&\quad + ((\bullet\bullet(\bullet\bullet))\bullet) - (((\bullet\bullet)\bullet\bullet)\bullet) + (((\bullet\bullet\bullet)\bullet)\bullet) - ((\bullet(\bullet\bullet))\bullet\bullet), \text{ and}\\
\partial(\bullet\bullet\bullet\bullet)^4 &:= ((\bullet\bullet)(\bullet\bullet)\bullet) - (\bullet\bullet((\bullet\bullet)\bullet)) + (\bullet(\bullet(\bullet\bullet)\bullet)) - (\bullet(\bullet(\bullet\bullet))\bullet)\\
&\quad + ((\bullet(\bullet\bullet\bullet))\bullet) - ((\bullet(\bullet\bullet)\bullet)\bullet) + (((\bullet\bullet\bullet)\bullet)\bullet) - (((\bullet\bullet)\bullet\bullet)\bullet).
\end{aligned}$$



FIGURE 2. Stasheff's associahedron K_5.



FIGURE 3. An closed edge path of length 8 in K_5 defining $\partial(\mu_5^1)$.

Let us indicate how we obtained the above formulas. We observed first that the degree-one subspace $\Gamma(\mu_2, \mu_3)(5)_1 \subset \Gamma(\mu_2, \mu_3)(5)$ is spanned by $\circ_i$-compositions of two μ_2's and one μ_3, i.e., in the bracket language, by nested bracketings of five •'s with two binary and one ternary bracket. These elements are in one-to-one correspondence with the edges of the 5th Stasheff associahedron K_5 shown in Figure 2, see [17, Section II.1.6].

Let $x_e \in \Gamma(\mu_2, \mu_3)(5)_1$ be the element indexed by an edge e of K_5. Clearly $\partial(x_e) = x_a + x_b$, where a, b are the endpoints of e and $x_a, x_b \in \Gamma(\mu_2)(5)_0$ the elements given by the nested bracketings of five •'s with three binary brackets corresponding to these endpoints. We concluded that the ∂-cycles in $\Gamma(\mu_2, \mu_3)(5)_1$ are generated by closed edge-paths of even length in K_5; the cycle corresponding to such a path $P = (e_1, e_2, \ldots, e_{2r})$ being

$$\sum_{1 \leq i \leq 2r} (-1)^{i+1} x_{e_i}.$$

Examples of these paths are provided by two adjacent pentagons in K_5 such as the ones shown in Figure 3. There are also three edge paths of length 4 given by the three square faces of K_5, but the corresponding cohomology classes have already been killed by the ∂-images of the compositions $\mu_3 \circ_i \mu_3$, $i = 1, 2, 3$. We showed that there are four linearly independent edge paths of length 8 that, together with the three squares, generate all edge paths of even length in K_5. The generators $\mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4$ correspond to these paths.

Also for $a \geq 6$ the 1-dimensional ∂-cycles in $\Gamma(\mu_2, \mu_3, \mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4)(a)_1$ are given by closed edge paths of even length in the associahedron K_a but one can show that they are all generated by the squares and the images of the paths as in Figure 3 under the face inclusions $K_5 \hookrightarrow K_a$. Therefore $(\Gamma(\mu_2, \mu_3, \mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4), \partial)$ is acyclic in degree 1, so $\mu_5^1, \mu_5^2, \mu_5^3, \mu_5^4$ are the only degree two generators of the minimal model of $\widetilde{\mathcal{A}ss}$.

From the above description of the minimal model of $\widetilde{\mathcal{A}ss}$ one easily gets the relevant part

$$C^1_{\widetilde{\mathcal{A}ss}}(A;A) \xrightarrow{\delta^1} C^2_{\widetilde{\mathcal{A}ss}}(A;A) \xrightarrow{\delta^2} C^3_{\widetilde{\mathcal{A}ss}}(A;A) \xrightarrow{\delta^3} C^4_{\widetilde{\mathcal{A}ss}}(A;A) \xrightarrow{\delta^4} \cdots$$

of the complex defining the deformation cohomology of an anti-associative algebra $A = (V, \mu)$. One has

- $C^1_{\widetilde{\mathcal{A}ss}}(A;A) = Hom(V,V)$
- $C^2_{\widetilde{\mathcal{A}ss}}(A;A) = Hom(V^{\otimes 2},V)$
- $C^3_{\widetilde{\mathcal{A}ss}}(A;A) = Hom(V^{\otimes 3},V)$, and
- $C^4_{\widetilde{\mathcal{A}ss}}(A;A) = Hom(V^{\otimes 5},V) \oplus Hom(V^{\otimes 5},V) \oplus Hom(V^{\otimes 5},V) \oplus Hom(V^{\otimes 5},V)$.

Observe that $C^p_{\widetilde{\mathcal{A}ss}}(A;A)_{\rm st} = C^p_{\widetilde{\mathcal{A}ss}}(A;A)$ for $p = 1,2,3$, while $C^4_{\widetilde{\mathcal{A}ss}}(A;A)$ consists of 5-linear maps. The differential δ^p agrees with $\delta^p_{\rm st}$ for $p = 1,2$ while, for $g \in C^3_{\widetilde{\mathcal{A}ss}}(A;A)$, one has

$$\delta^3(g) = (\delta^3_1(g), \delta^3_2(g), \delta^3_3(g), \delta^3_4(g)),$$

where

$$\begin{aligned}
\delta^3_1(g)(a,b,c,d,e) :=\ & af(b,c,de) - f(a,b,c(de)) + (ab)f(c,d,e) - f(ab,cd,e) \\
& + f(ab,c,d)e - f((ab)c,d,e) + f(a,b,c)(de) - f(a,bc,de), \\
\delta^3_2(g)(a,b,c,d,e) :=\ & f((ab)c,d,e) - f(ab,c,d)e + f(a,b,cd)e - f(a,b(cd),e) \\
& + af(b,cd,e) - f(a,b,(cd)e) + (ab)f(c,d,e) - f(ab,c,de), \\
\delta^3_3(g)(a,b,c,d,e) :=\ & f(a,bc,de) - af(bc,d,e) + f(a,(bc)d,e) - a(f(b,c,d)e) \\
& + f(a,b,cd)e - f(ab,c,d)e + (f(a,b,c)d)e - f(a(bc),d,e), \text{ and} \\
\delta^3_4(g)(a,b,c,d,e) :=\ & f(ab,cd,e) - f(a,b,(cd)e) + af(b,cd,e) - f(a,b(cd),e) \\
& + (af(b,c,d))e - f(a,bc,d)e + (f(a,b,c)d)e - f(ab,c,d)e,
\end{aligned}$$

for $a,b,c,d,e \in V$. The following proposition follows from [12, Section 4].

Proposition 19. *The cohomology $H^*_{\widetilde{\mathcal{A}ss}}(A;A)$ governs deformations of anti-associative algebras. This means that $H^2_{\widetilde{\mathcal{A}ss}}(A;A)$ parametrizes isomorphism classes of infinitesimal deformations and $H^3_{\widetilde{\mathcal{A}ss}}(A;A)$ contains obstructions to extensions of partial deformations.*

6. Free partially associative n-algebras

In this section we describe free partially associative algebras in the situation when the corresponding operad is Koszul. We thus fill in the gap in the proof of [7, Theorem 13] and extend Gnedbaye's description of free $p\mathcal{A}ss^n_d$-algebras to all cases when d and n have the same parity.

Let $p\mathcal{A}ss^n_d(V)$ be the free $p\mathcal{A}ss^n_d$-algebra generated by a graded vector space V. It obviously decomposes as

$$p\mathcal{A}ss^n_d(V) = \bigoplus_{l \geq 0} p\mathcal{A}ss^n_d(V)_l,$$

where $p\mathcal{A}ss^n_d(V)_l \subset p\mathcal{A}ss^n_d(V)$ is the subspace generated by elements obtained by applying the structure n-ary multiplication μ to elements of V l-times. For instance, $p\mathcal{A}ss^n_d(V)_0 \cong V$ and $p\mathcal{A}ss^n_d(V)_1 \cong V^{\otimes n}$.

Denote by $\mathcal{T}^n_l$, $l \geq 1$, the set of planar directed (= rooted) trees with $l(n-1)+1$ leaves whose vertices have precisely n incoming edges (see [16, Section 4] or [17, II.1.5] for terminology). We extend the definition to $l = 0$ by putting $\mathcal{T}^n_0 := \{ \mid \}$, the one-point set consisting of the exceptional tree with one leg and no internal vertex. Clearly, each tree in $\mathcal{T}^n_l$ has exactly l vertices. For each l there is a natural epimorphism

$$\omega : \mathcal{T}^n_l \times V^{\otimes l(n-1)+1} \twoheadrightarrow p\mathcal{A}ss^n_d(V)_l \tag{13}$$

given by interpreting the trees in $\mathcal{T}^n_l$ as the 'pasting schemes' for the iterated multiplication μ. More precisely, if $T \in \mathcal{T}^n_l$ and $v_1, \ldots, v_{l(n-1)+1} \in V$, then

$$\omega(T \times (v_1, \ldots, v_{l(n-1)+1})) \in p\mathcal{A}ss^n_d(V)_l$$

is obtained by decorating the vertices of T by μ, the leaves of T by elements $v_1, \ldots, v_{l(n-1)+1}$, and performing the indicated composition, observing the Koszul sign rule in the nontrivially graded cases.

Let $\mathcal{S}^n_l$ be the subset of $\mathcal{T}^n_l$ of trees having the property that the first incoming edge of each vertex is a leaf. Therefore $\mathcal{S}^n_0 = \mathcal{T}^n_0 = \{ \mid \}$, $\mathcal{S}^n_1$ is the one-point set consisting of the n-corolla



and $\mathcal{S}^n_2$ has $n-1$ elements



, $2 \leq i \leq n$.

It is clear that, for $l \geq 3$, $\mathcal{S}_l^n$ consists of trees of the form

(14)



where $S_i \in \mathcal{S}_{l_i}^n$ for $2 \leq i \leq n$ and $l_2 + \cdots + l_n = l - 1$. The main result of this section is

Theorem 20. *Assume that n and d are of the same parity, so the operad $p\mathcal{A}ss_d^n$ is Koszul by Theorem 9. Then the restriction (denoted by the same symbol)*

$$\omega : \mathcal{S}_l^n \times V^{\otimes l(n-1)+1} \longrightarrow p\mathcal{A}ss_d^n(V)_l \tag{15}$$

of the epimorphism (13) is an isomorphism, for each $l \geq 0$.

Proof. Axiom (2) for partially associative algebras implies that each iterated multiplication in $p\mathcal{A}ss_d^n(V)_l$ can be brought into a linear combination of multiplications with the pasting schemes in $\mathcal{S}_l^n$, i.e. that the map (15) is an epimorphism. Let us prove this statement by induction.

There is nothing to prove for $l = 0, 1$. Assume that we have established the claim for all $0 \leq l \leq k$, $k \geq 1$, and prove it for $l = k$. Let μ_T be an iterated multiplication with the pasting scheme $T \in \mathcal{T}_k^n$. There are two possibilities. The *first case:* the tree T is of the form

(16)



for some $T_i \in \mathcal{T}_{l_i}^n$, $2 \leq i \leq n$, with $l_2 + \cdots + l_n = k - 1$. Then

$$\mu_T = \mu(1\!\!1 \otimes \mu_{T_2} \otimes \cdots \otimes \mu_{T_n}),$$

where μ_{T_i} denotes the iterated multiplication with the pasting scheme T_i. By induction, each $\mu_{T_2}, \ldots, \mu_{T_n}$ is a linear combination of iterated multiplications whose pasting schemes belong to the subsets $\mathcal{S}_{l_2}^n, \ldots, \mathcal{S}_{l_n}^n$, respectively. The observation that the tree (16) belongs to $\mathcal{S}_k^n$ if $T_i \in \mathcal{S}_{l_i}^n$ for each $2 \leq i \leq n$ completes the induction step for this case.

In the *second case,* the tree T has the form



where $T_i \in \mathcal{T}_{l_i}^n$ for $1 \leq i \leq n$, $l_1 + \cdots + l_n = k - 1$ and $l_1 \geq 1$. Now

$$\mu_T = \mu(\mu_{T_1} \otimes \cdots \otimes \mu_{T_n})$$

and we may assume, by induction, that $T_i \in \mathcal{S}^n_{l_i}$ for each $1 \leq i \leq n$. In particular, T_1 is as in (14) with $S_j \in \mathcal{S}^n_{l'_j}$, $2 \leq j \leq n$ such that $l'_2 + \cdots + l'_n = l_1 - 1$, thus

$$\mu_T = \mu(\mu \otimes 1\!\!1^{\otimes n-1})(1\!\!1 \otimes \mu_{S_2} \otimes \cdots \otimes \mu_{S_n} \otimes \mu_{T_2} \otimes \cdots \otimes \mu_{T_n}).$$

By (2), one may replace the factor $\mu(\mu \otimes 1\!\!1^{\otimes n-1})$ by the linear combination

$$-\sum_{i=2}^{n} (-1)^{(i+1)(n-1)} \mu \left(1\!\!1^{\otimes i-1} \otimes \mu \otimes 1\!\!1^{\otimes n-i}\right)$$

which brings μ_T also in the second case to the desired form and finishes the induction step.

To prove that the map (15) it is an isomorphism, it suffices now to compare the dimensions of $\mathcal{S}^n_l \times V^{\otimes l(n-1)+1}$ and $p\mathcal{A}ss^n_d(V)_l$. It follows from the description [17, Proposition II.1.25] of the free operad algebra that, for each $l \geq 0$,

$$p\mathcal{A}ss^n_d(V)_l \cong p\mathcal{A}ss^n_d(l(n-1)+1) \otimes \Sigma_{l(n-1)+1} V^{\otimes l(n-1)+1}.$$

Theorem 20 will thus be established if we prove that

$$S^n_l := \mathrm{card}(\mathcal{S}^n_l) \text{ equals } A^n_l := \frac{1}{(l(n-1)+1)!} \dim(p\mathcal{A}ss^n_d(l(n-1)+1)),$$

for each $l \geq 0$. It easily follows from (14) that the sequence $\{S^n_l\}_{l \geq 0}$ is determined by the recursion $S^n_0 = 1$ and

$$S^n_l := \sum_{\substack{0 \leq l_2, \cdots, l_n \leq l-1 \\ l_2 + \cdots + l_{n-1} = l-1}} S^n_{l_2} \cdots S^n_{l_n} \text{ for } l \geq 1. \tag{17}$$

In Proposition 21 below, which is of independent interest, we prove that the sequence $\{A^n_l\}_{l \geq 0}$ satisfies the same recursion. This finishes the proof. □

Recursion (17) appeared, with $p_l^{<n-1>}$ in place of S^n_l, in [7, Section 3.4]. Theorem 20 gives a realization of free $p\mathcal{A}ss^n_d$-algebras in the Koszul case ($n \equiv d$ mod 2) by putting

$$p\mathcal{A}ss^n_d(V) := \bigoplus_{l \geq 0} \mathcal{S}^n_l \times V^{\otimes l(n-1)+1}.$$

We leave as an exercise to describe the structure n-ary multiplication of $p\mathcal{A}ss^n_d(V)$ in this language, see [7].

Proposition 21. *The Poincaré series of the operad $p\mathcal{A}ss^n_d$ is, in the Koszul case (with n and d of the same parity), given by*

$$g_{p\mathcal{A}ss^n_d}(t) = \sum_{l \geq 0} (-1)^{ln} A^n_l t^{l(n-1)+1} \tag{18}$$

where the coefficients $\{A^n_l\}_{l \geq 0}$ are defined recursively by $A^n_0 := 1$ and

$$A^n_l := \sum_{\substack{0 \leq l_2, \cdots, l_n \leq l-1 \\ l_2 + \cdots + l_{n-1} = l-1}} A^n_{l_2} \cdots A^n_{l_n} \text{ for } l \geq 1. \tag{19}$$

Proof. One can easily check that the recursive definition (19) of the coefficients of $f(t) := g_{p\mathcal{A}ss^n_d}(t)$ is equivalent to the functional equation

$$f(t) = t\left(1 + (-1)^n f(t)^{n-1}\right)$$

which in turn immediately implies that $f(t)$ is the unique formal solution of

$$g_{t\mathcal{A}ss^n_{-d+n-2}}(-f(-t)) = t,$$

where the Poincaré series $g_{t\mathcal{A}ss^n_{-d+n-2}}(t)$ is as in the first line of (6) because $-d+n-2$ is even. Since we are in the Koszul case, the above display means, by Theorem 12, that $f(t)$ is the Poincaré series of $(t\mathcal{A}ss^n_{-d+n-2})^! = p\mathcal{A}ss^n_d$. This proves the proposition. □

The description of the Poincaré series of $p\mathcal{A}ss^n_d$ for n and d of the same arity given in Proposition 21 implies that the Poincaré series of $\widetilde{p\mathcal{A}ss^n_d}$ for d odd equals

$$g_{\widetilde{p\mathcal{A}ss^n_d}}(t) = \sum_{l\geq 0}(-1)^l A^n_l t^{l(n-1)+1},$$

with $\{A^n_l\}_{l\geq 0}$ having the meaning as in (18).

Example 22. Using `Matematica`, we calculated initial values of the series $\{A^3_l\}_{l\geq 0}$ as $A^3_0 = 1$, $A^3_1 = 1$, $A^3_2 = 2$, $A^3_3 = 5$, $A^3_4 = 14$, $A^3_5 = 42$, $A^3_6 = 132$, $A^3_7 = 429$, $A^3_8 = 1\ 430$, $A^3_9 = 4\ 862$, $A^3_{10} = 16\ 796$, &c.

Remark 23. In the non-Koszul case (n and d of different parity), the map (15) of Theorem 20, while always being an epimorphism, need not be a monomorphism. This means that there may be "unexpected relations" in the free algebra $p\mathcal{A}ss^n_d(V)$ and Gnedbaye's description of free partially associative n-algebras is no longer true.

For instance, while S^3_3 (the dimension of $\mathcal{S}^n_3$ for $n = 3$) equals 5, the dimension of $p\mathcal{A}ss^3_0(7)$ equals $7! \cdot 4$, so $p\mathcal{A}ss^3_0(V)_3$ has one copy $V^{\otimes 7}$ less than $\mathcal{S}^3_3 \times V^{\otimes 7}$. Therefore the map (15) has, for $n = 3$, $d = 0$ and $l = 3$, a nontrivial kernel. The Poincaré series of $p\mathcal{A}ss^3_0$ is was calculated in [8] as

$$g_{p\mathcal{A}ss^3_0}(t) = t + t^3 + 2t^5 + 4t^7 + 5t^9 + 6t^{11} + 7t^{13} + \cdots .$$

7. Open problems

The first question which our paper leaves open is the Koszulity of the operads $t\mathcal{A}ss^n_d$ with d odd and $n \geq 8$. The method used in the proof of Proposition 17 does not apply to these cases and indeed, our numerical tests mentioned in Remark 18 suggest that it may happen that all coefficients in the formal inverse of $t - t^n + t^{2n-1}$ are non-negative.

Even if this happens, it would not necessarily mean that the operad $t\mathcal{A}ss^n_d$ is Koszul, only that subtler methods must be applied to that case. For instance, one may try to compare the coefficients of this formal inverse to the dimensions of the components of the dual operad $(t\mathcal{A}ss^n_d)^!$.

Understanding these components is, of course, equivalent to finding a basis for the free partially associative algebras in the non-Koszul cases. This problem was solved, in [8], for free $p\mathcal{A}ss^3_0$-algebras, for $n \geq 4$ it remains open.

The last problem we want to formulate here is to find more about the minimal model of the anti-associative operads $\widetilde{\mathcal{A}ss}$, or even to describe it completely. As far as we know, no complete description of the minimal model of a non-Koszul operad is known. Since $\widetilde{\mathcal{A}ss}$ is the simplest non-Koszul operad, it is the first obvious candidate to attack. A related task is to find as much as information about minimal models of the remaining non-Koszul n-ary operads as possible.

References

[1] H. Ataguema and A. Makhlouf. Notes on cohomologies of ternary algebras of associative type. Preprint `arXiv:0812.0707`, December 2008.

[2] V. Dotsenko and A. Khoroshkin. Gröbner bases for operads. Preprint `arXiv:0812.4069`, December 2008.

[3] T.F. Fox. An introduction to algebraic deformation theory. *Journal of Pure and Appl. Algebra*, 84:17–41, 1993.

[4] T.F. Fox and M. Markl. Distributive laws, bialgebras, and cohomology. In J.-L. Loday, J.D. Stasheff, and A.A. Voronov, editors, *Operads: Proceedings of Renaissance Conferences*, volume 202 of *Contemporary Math.*, pages 167–205. Amer. Math. Soc., 1997.

[5] B. Fresse. Koszul duality of operads and homology of partition posets. *Contemp. Math.*, 346:115–215, 2004.

[6] V. Ginzburg and M.M. Kapranov. Koszul duality for operads. *Duke Math. J.*, 76(1):203–272, 1994.

[7] A.V. Gnedbaye. Opérades des algèbres $(k+1)$-aires. In J.L. Loday, J.D Stasheff, and A.A. Voronov, editors, *Operads: Proceedings of Renaissance Conferences*, volume 202 of *Contemporary Math.*, pages 83–113. Amer. Math. Soc., 1996.

[8] N. Goze and E. Remm. n-ary associative algebras, cohomology, free algebras and coalgebras. Preprint `arXiv:0803.0553`, March 2008.

[9] P. Hanlon and M.L. Wachs. On Lie k-algebras. *Adv. in Math.*, 113:206–236, 1995.

[10] E. Hoffbeck. A Poincaré-Birkhoff-Witt criterion for Koszul operads. Preprint `arXiv:0709.2286`, September 2007.

[11] M. Markl. A cohomology theory for $A(m)$-algebras and applications. *J. Pure Appl. Algebra*, 83:141–175, 1992.

[12] M. Markl. Cotangent cohomology of a category and deformations. *J. Pure Appl. Algebra*, 113(2):195–218, 1996.

[13] M. Markl. Models for operads. *Comm. Algebra*, 24(4):1471–1500, 1996.

[14] M. Markl. Intrinsic brackets and the L_∞-deformation theory of bialgebras. Preprint `math.AT/0411456`, November 2004.

[15] M. Markl. A resolution (minimal model) of the PROP for bialgebras. *J. Pure Appl. Algebra*, 205(2):341–374, 2006.

[16] M. Markl. *Handbook of Algebra*, volume 5, chapter Operads and PROPs, pages 87–140. Elsevier, 2008.

[17] M. Markl, S. Shnider, and J. D. Stasheff. *Operads in Algebra, Topology and Physics*, volume 96 of *Mathematical Surveys and Monographs.* American Mathematical Society, Providence, Rhode Island, 2002.

MATHEMATICAL INSTITUTE OF THE ACADEMY, ŽITNÁ 25, 115 67 PRAGUE 1, THE CZECH REPUBLIC

E-mail address: `markl@math.cas.cz`

Laboratoire de Mathématiques et Applications, Université de Haute Alsace, Faculté des Sciences et Techniques, 4, rue des Frères Lumière, 68093 Mulhouse cedex, France.

E-mail address: `Elisabeth.Remm@uha.fr`